

1 **Genetic structure of a patchily distributed philopatric migrant: implications for**
2 **management and conservation**

3 Anastasios Bounas^{1,2*}, Dimitris Tsaparis³, Ron Efrat⁴, Gradimir Gradev⁵, Marco Gustin⁶,
4 Kresimir Mikulic⁷, Airam Rodríguez⁸, Maurizio Sarà⁹, Georgios Kotoulas³, Konstantinos
5 Sotiropoulos¹

6 ¹*Molecular Ecology and Conservation Genetics Lab, Department of Biological Applications*
7 *and Technology, University of Ioannina, 45110, Ioannina, Greece.*

8 ²*Hellenic Ornithological Society – BirdLife Greece, Themistokleous 80, 10681 Athens, Greece.*

9 ³*Institute of Marine Biology, Biotechnology and Aquaculture, Hellenic Centre for Marine*
10 *Research, Heraklion, 71500, Greece.*

11 ⁴*Mitrani Department of Desert Ecology, The Jacob Blaustein Institutes for Desert Research,*
12 *Ben-Gurion University of the Negev, IL-8499000, Israel.*

13 ⁵*Green Balkans NGO, 1 Skopie Street, 4004, Plovdiv, Bulgaria.*

14 ⁶*LIPU (Lega Italiana Protezione Uccelli) - BirdLife Italia, Conservation Department, Via Udine*
15 *3, I-43100, Parma, Italy.*

16 ⁷*Association BIOM – BirdLife Croatia, Preradoviceva 34, 10000, Zagreb, Croatia.*

17 ⁸*Department of Evolutionary Ecology, Estación Biológica de Doñana (CSIC), Avda. Américo*
18 *Vespucio 26, 41092, Seville, Spain.*

19 ⁹*Section of Animal Biology, Department STEBICEF, University of Palermo, Via Archirafi 18,*
20 *90123 Palermo, Italy.*

21

22 *Corresponding author: abounas@cc.uoi.gr, tel: +306973573235

23 Running Title: Conservation Genetics of the Lesser Kestrel

24

Significant demographic fluctuations can have major genetic consequences in wild populations. The Lesser Kestrel (*Falco naumanni*) has suffered from both population declines and range fragmentation during the second half of the 20th century. In this study we analysed multilocus microsatellite data to assess the genetic structure of the species. Our analysis revealed significant genetic structuring of Lesser Kestrel populations, not only at cross-continental scale, but also regionally within Central and Eastern (CE) Mediterranean. We detected signs of genetic bottlenecks in some of the peripheral populations coupled with small effective population sizes. Values of genetic differentiation among the largest populations were low, albeit significant, whereas the small peripheral CE Mediterranean populations showed higher levels of differentiation from all other populations. Gene flow levels were relatively low among the discontinuously distributed populations of the CE Mediterranean region. We argue that the observed spatial genetic structure can be attributed at some level to the past demographic decline experienced by the species. Finally, we identify management units in the region, and inform the design of conservation actions aiming at the increase of population sizes and dispersal rates among peripheral populations.

Keywords: *Falco naumanni* -Genetic Diversity-Lesser Kestrel-Management Units-Mediterranean-Microsatellites-Migration Rates

48 INTRODUCTION

49 In many animal species, the patterns of genetic differentiation and gene flow are highly
 50 influenced by the geographical characteristics of their habitats as well as their migratory
 51 behavior (Willoughby *et al.*, 2017). Birds, and specifically raptors, can show long-distance
 52 migratory behavior and also exhibit natal and breeding site fidelity. As a general pattern,
 53 migratory populations of raptors have been found to have weaker genetic structure and
 54 higher genetic diversity than resident populations (Miller *et al.*, 2012; Willoughby *et al.*,
 55 2017). For example some individuals might migrate through a different route upon their
 56 return on the breeding grounds, and consequently disperse and breed far from their natal
 57 site (Garcia *et al.*, 2011). On the other hand, philopatry reduces or even inhibits gene flow
 58 among populations leading to increased genetic differentiation. Patchily distributed and
 59 locally isolated populations are susceptible to a greater influence of genetic drift which may
 60 result in a decrease in genetic diversity and fitness, thus compromising a species' ability to
 61 adapt to a changing environment (Amos & Balmford, 2001). Genetic drift and inbreeding are
 62 expected to be stronger in peripheral populations relative to core ones, due to their small
 63 population sizes and low immigration rates (Vucetich & Waite, 2003; Hanski & Gaggiotti,
 64 2004). Therefore, the assessment of genetic structure and the identification of its underlying
 65 processes become essential tasks providing valuable information towards the design and
 66 implementation of effective conservation strategies. For example, identification of
 67 management units (MUs, Moritz, 1994; Taylor & Dizon, 1999; Palsbøll, Berube & Allendorf,
 68 2007) would be central to delineate populations for monitoring and thus aid their short-term
 69 management.

70 The Lesser Kestrel (*Falco naumanni*, Fleischer, 1818) is a small migratory falcon breeding
 71 from the Mediterranean basin across Middle East and Central Asia to Mongolia and China,
 72 and wintering in sub-Saharan Africa (Cramp & Simmons, 1980). The species underwent rapid

declines throughout its European range in the early 1960s mainly as a result of agricultural intensification and subsequent habitat degradation and land use changes (Iñigo & Barov, 2010). In Central-Eastern Mediterranean region (hereafter CE Mediterranean) and especially in the Balkan Peninsula, the decline was dramatic, leading to local extinctions and consequently to significant range contraction and fragmentation (Iñigo & Barov, 2010). Currently, the species shows a patchy distribution and is considered to have a “depleted” status in the region (BirdLife International, 2017a). The two largest (core) populations are located in the Apulia-Basilicata area of southern Italy and in Central Greece, still holding several thousand pairs (BirdLife International, 2017a). Peripheral populations of smaller size, still exist in the Former Yugoslav Republic of Macedonia (Uzunova & Lisichanets, 2016), in Sicily (Sarà, 2004), the European part of Turkey (Kmetova *et al.*, 2012) and throughout continental Greece as well as on some of the Greek islands (Legakis & Maragou, 2009). Finally, a small geographically isolated population is located in Croatia, at the northernmost edge of the species European distribution (Mikulic *et al.*, 2013). Certain conservation actions, mainly implemented in Western Europe, have led to a stabilization and slightly positive population trend and subsequently the down-listing of the species from Vulnerable to Least Concern (BirdLife International, 2017b).

Several previous studies have examined the genetic structure of the species and the underlying processes, at both continental (Wink, Sauer-Gurth & Pepler, 2004; Alcaide *et al.*, 2008a; Alcaide *et al.*, 2008b) and finer spatial scales (Ortego *et al.*, 2008a; Ortego *et al.*, 2008b; Alcaide *et al.*, 2009; Di Maggio *et al.*, 2014). It has been proposed that Asian Lesser Kestrels are considerably differentiated from European conspecifics, whereas populations across the Western Palearctic seem to follow an isolation by distance pattern while maintaining high levels of genetic diversity (Wink *et al.*, 2004; Alcaide *et al.*, 2008b). On a regional scale though, it seems that the species' philopatry does not lead to fine-scale genetic structuring (Alcaide *et al.*, 2009); nevertheless, population structure could emerge,

depending on the size and the degree of spatial isolation and the levels of gene flow among populations (Ortego *et al.*, 2008b). Such restricted gene flow patterns do not necessarily result from physical barriers hindering dispersal, especially in efficient dispersers such as the Lesser Kestrel, but can be attributed also to their philopatric behaviour. Indeed, the species shows high philopatry, with adult birds returning to breed close to their previous breeding territories (Negro, Hiraldo & Donázar, 1997), however, juveniles disperse more and in greater distances (Serrano & Tella, 2003; Bounas *et al.*, 2016a). At least at a continental scale, Lesser Kestrels show strong migratory connectivity, i.e. there is a spatial segregation of breeding populations at the wintering range: European populations winter in the Sahel, while Asian populations winter in east and South Africa (Wink *et al.*, 2004; Rodriguez *et al.*, 2009)

Herein, we examine the genetic structure of the Lesser Kestrel, 1) across the broader species range and 2) within the CE Mediterranean. We aim to identify patterns of genetic variation and gene flow among populations as well as their underlying processes. Such information could be of crucial importance for conservation programs to identify the need of local scale conservation actions and inform their design.

MATERIAL AND METHODS

POPULATION SAMPLING AND DNA EXTRACTION

Samples were obtained from 12 breeding sites in CE Mediterranean, where the species shows a fragmented distribution as well as from Mongolia, Israel and two sites from Spain. Individuals sampled in Bulgaria originated from Spain and were used for the reinforcement of the species, thus treated as if they were sampled in Extremadura (ES/BG; Fig. 1). A total of 295 individuals were sampled during four consecutive breeding seasons (2013 - 2016) with the exception of Spanish samples (SES) that were collected in 2007. Birds were caught by

hand in the nest or using mist nets or spring traps close to the nest. To minimize biases associated with relatedness only a single fledgling per brood was sampled. Two drops of blood (~50µl) were obtained from each individual by leg-pricking and immediately stored in blood storage cards (NucleoCards®) at room temperature until DNA extraction. DNA was extracted using the NucleoSpin Tissue kit (Macherey-Nagel) following the manufacturer's protocol.

MICROSATELLITE AMPLIFICATION AND GENOTYPING

Each individual was genotyped at a total of 18 microsatellite loci. Seven loci were originally isolated from the Peregrine Falcon *Falco peregrinus* (Nesje *et al.*, 2000; Alcaide *et al.*, 2008a), whereas 11 were developed specifically for the Lesser Kestrel (Ortego *et al.*, 2007; Padilla *et al.*, 2009). Details on loci properties and primers used for their amplification are presented in Supporting information, Table S1. All loci were amplified in five multiplex reactions using forward 5'-fluorescent-labelled primers and the KAPA2G Fast Multiplex PCR Kit (Kapa Biosystems). Each 12.5 µl reaction contained 2pM of each primer and 1x KAPA2G Mix and was carried out using the following profile: an initial denaturation step of 3 min at 95 °C, 30 cycles of 15 s at 95°C, 30 s at 60 °C, 30 s at 72 °C, with a final extension step of 10 min at 72 °C. PCR products were separated and visualized using an ABI 3730xl capillary sequencer (Applied Biosystems) and genotypes were scored by eye with STRand v.2.4.59 (Toonen & Hughes, 2001). Randomization of samples was employed throughout lab processes to avoid any plate/gel specific errors that might lead in population specific biases (Meirmans, 2015). In addition, a subset of 50 individuals was re-genotyped to quantify error rates due to allelic dropout or genotyping errors but no inconsistencies were detected. We used the package "MsatAllele" (Alberto, 2009) in R 3.2.2 (R Core Team, 2015) to allocate alleles to their respective size classes. Genotyping errors, due to null alleles and stuttering, were examined for all loci and sampled populations using MICROCHECKER (Van Oosterhout *et al.*, 2004).

Two loci (Fnd1.2 and Fnd2.1) were omitted from further analyses due to the presence of null alleles and extensive stuttering respectively, and thus the full analysis was based on the remaining 16 microsatellite loci.

GENETIC ANALYSES

To evaluate the genetic variability in each population, standard genetic diversity indices (A: number of alleles, H_O : observed and H_E : expected heterozygosity) were calculated using the program GENALEX v.6.5 (Peakall & Smouse, 2012). Rarefied Private Allelic richness (π) estimates were produced using HP-RARE (Kalinowski, 2005). Allelic richness (A_R) corrected for different sample sizes was calculated using FSTAT 2.9.3.2 (Goudet, 2002). The same software was used to calculate the inbreeding coefficient (F_{IS}), test for deviations from Hardy-Weinberg proportions at each locus and sampled population, as well as for linkage disequilibrium (LD) using 1000 randomizations and adjusting significance for multiple comparisons (adjusted P value < 0.0003). In addition we performed a Chi-Square test and Fisher's method to confirm the Hardy Weinberg (HW) equilibrium results.

The software Arlequin 3.5.1.3 (Excoffier & Lischer, 2010) was used to calculate F_{ST} values between all pairs of sampled populations and test them for statistical significance using 10000 permutations. In addition to F_{ST} , we also calculated Jost's D (D_{est}) as an unbiased estimator of differentiation that performs better than other relatives, in cases of markers with different number of alleles (Gerlach *et al.*, 2010) as is the case in this study (see Supporting information, Table S1). Pairwise D values (mean D_{est}) among populations were calculated with the R-package "DEMEtics" (Gerlach *et al.*, 2010) and statistical significance was tested using 1,000 bootstrap iterations. For both estimators, p-values were adjusted for multiple comparisons after the B-H method (Benjamini & Hochberg, 1995). We used IBDWS v.3.23 (Jensen, Bohonak & Kelley, 2005) in order to obtain any statistically significant associations between pairwise genetic ($F_{ST}/[1-F_{ST}]$) and linear geographic (log km) distance

matrices using 30,000 randomizations. The analysis was performed both on the full and the reduced (i.e. CE Mediterranean) datasets.

To evaluate the genetic population structure, the Bayesian clustering software STRUCTURE 2.3.4 (Pritchard, Stephens & Donnelly, 2000) was used to infer the number of genetically homogeneous clusters present in the dataset. The analysis was conducted both on the full dataset as well as only on the populations of CE Mediterranean region. For both analyses we assumed the admixture ancestry model and correlated allele frequencies (Falush, Stephens & Pritchard, 2003), using sampling location as prior information (LOCPRIOR model; Hubisz *et al.*, 2009) as it is deemed to be more sensitive at inferring population structure at lower levels of divergence, which is expected in good dispersers such as birds. Runs were set with a burn-in period of 2×10^5 iterations followed by 10^6 MCMC steps with 20 replicates for each K value (1 to 15 for the full dataset and 1 to 12 for CE Mediterranean). STRUCTURE runs were implemented on a beowulf cluster using PARASTRUCTURE Perl script (Lagnel, 2015) and STRUCTURE plots were constructed using DISTRUCT (Rosenberg, 2004). The most likely value of genetic clusters, K, was evaluated following the ΔK method (Evanno, Regnaut & Goudet, 2005) implemented in STRUCTURE HARVESTER (Earl & Vonholdt, 2012), as well as by calculating the posterior probability for each K. We present all cases with high probability that warrant a biological interpretation (Meirmans, 2015) and are supported by both STRUCTURE runs.

Finally, the breeding populations of CE Mediterranean were grouped according to Structure results as well as F_{st} and D_{est} calculations, and the directional contemporary gene flow and its relative magnitude among them was estimated using the divMigrate function (Sundqvist *et al.* 2016) in the R-package “diveRsity” (Keenan *et al.* 2013). The method provides a relative (to within the analysis) migration network graph aiming to visualize the gene flow patterns among populations, with the resulting metric representing a proportion

of the gene flow among areas, scaled to the largest magnitude estimated. Populations are represented as nodes and the properties of the lines connecting them are based on the relative strength of gene flow. Populations that exhibit strong gene exchange between them but weak gene flow with others tend to cluster closely together, reflecting patterns of genetic structure. The method is described in detail in Sundqvist *et al.* (2016). We used N_M (Alcala, Goudet & Vuilleumier, 2014) as a measure of genetic distance, and tested whether gene flow among populations was asymmetrical using 10000 bootstrap iterations.

Signs of bottlenecks were evaluated in grouped populations using three approaches. First we calculated M , a ratio based on the number of alleles to the allelic size range (Garza & Williamson, 2001) in Arlequin. M will be smaller in populations that have suffered a decline than in populations that are in mutation-drift equilibrium. A test for heterozygosity excess was performed in BOTTLENECK (Piry, Luikart & Cornuet, 1999) using the Wilcoxon signed rank test running 1000 iterations and using the two-phase model (TPM). As the microsatellites we used are either of dinucleotide perfect repeats or of imperfect repeats, both of which may tend toward the infinite allele model (IAM; Cornuet and Luikart, 1996), we fixed the proportions of the TPM in favour of the IAM (Cristescu *et al.*, 2010) including 20% of the stepwise mutation model (SMM) and 80% of IAM. We finally tested for a mode-shift distortion using a graphical approach, by plotting the number of alleles in 10 allele frequency classes with 0.1 intervals, according to Luikart *et al.* (1998). In a population at equilibrium, alleles with frequencies in the first class (<0.1) are expected to be more numerous than those belonging to the second class, therefore allele frequencies present a characteristic L-shaped distribution. Plots were examined for mode-shift distortions that would be consistent with a bottlenecked population (Luikart *et al.*, 1998; Cristescu *et al.*, 2010). Plots were produced in R 3.2.2 and allele frequency calculations were performed using the R package “Gstudio” (Dyer, 2014). Finally, the effective population size (N_E) along with 95% confidence intervals (CIs) were estimated using the bias-corrected version of the

method based on linkage disequilibrium (Hill, 1981; Waples, 2006) as implemented in NeESTIMATOR v.2.01 (Do *et al.*, 2014). N_E estimates were obtained for all but the core populations (ES, APU, CNG) and Mongolia, as it is very difficult to obtain reliable estimates for large populations using this method (Waples & Do, 2010).

RESULTS

All microsatellite markers were found to be polymorphic across populations and the number of alleles per locus ranged from four (loci Fnd1.4, Fp86-2, Fp89) to 38 (locus Fnd1.6; Supporting information, Table S1).

GENETIC DIVERSITY

We did not detect any differences in the mean allelic richness across populations. The average H_O was 0.64, H_O and H_E were similar and F_{IS} was not significant in all cases (Table 1). Private alleles were present in all sampled populations with the exception of Croatia (CRO), Komotini (KAL) and one subpopulation in Central Greece (TRI). Generally, populations did not deviate from Hardy-Weinberg proportions but some deviations of individual loci were detected: four populations (CRO, LIM, ISR, MON) showed deviations at two, one, one and two out of the 16 loci respectively (see Supporting information, Table S4). Since these loci did not show consistent deviations across all populations, we included them in subsequent analyses attributing this disequilibrium in processes specific to those populations. No LD was detected between any of the loci across all populations.

POPULATION STRUCTURE

Pairwise F_{ST} and D_{est} values were highly correlated (Pearson's correlation: $r = 0.95$, $P < 0.001$). Estimates of both F_{ST} and D_{est} (Table 2) varied between population pairs (F_{ST} : 0.003 – 0.06; D_{est} : 0.02 – 0.19), with all the core populations of Europe (Spain, Apulia, Central Greece) showing low differentiation among them. On the other hand, Mongolia and Israel seem

differentiated from all others, while the small peripheral CE Mediterranean populations showed higher levels of differentiation from all other populations. This was particularly evident for the populations of Croatia and Limnos where the highest values of D_{est} were recorded when compared to all other populations (Table 2). Mantel tests showed that genetic distance is not significantly correlated with geographical distance across all sampled populations ($r = 0.33$, $P = 0.1$, Supporting information, Figure S1). Conversely, genetic divergence of CE Mediterranean populations correlated significantly with the geographical distance among them ($r = 0.57$, $P = 0.01$, Supporting information, Figure S1).

The Bayesian clustering method implemented in STRUCTURE suggested the presence of population structuring. When all individuals were modeled, the ΔK -method suggested two clusters ($K = 2$) as the most likely population structure (although the posterior probability was higher for $K = 3$; Supporting information, Figure S2, Table S2). For $K = 2$ the admixture model indicated two gene pools with all European populations showing high membership coefficients in the first cluster, and the MON and LIM populations belonging to the second (Fig. 2). This second gene pool seems to be present in all populations in Northern and Central Greece (CNG) whereas absent from all other western populations. Lesser Kestrels from Israel (ISR) were found to be highly admixed, exhibiting a mixed ancestry from both clusters (Fig. 2). For $K = 3$, the model adds another gene pool that is mostly represented by the Trans-Adriatic populations of Croatia (CRO), Apulia (APU) and Sicily (SIC) whereas Israel (ISR) still seems to be of mixed ancestry. All subpopulations within Spain (SES, ES/BG) and Central Greece (TRI, LAR, VOL) showed identical admixture proportions among them, implying no further substructure (Fig. 2).

When individuals from the CE Mediterranean were modeled separately, the ΔK -method suggested $K = 2$ as the optimal number of clusters, whereas posterior probability estimates suggested the presence of four clusters (Supporting information, Figure S3, Table S3). The

populations of Croatia (CRO) and Limnos (LIM) constitute two different genetic clusters, the one present in Limnos (LIM) extending throughout Greek populations from Northern to Central Greece (CNG), while two other clusters are present in all other populations in different proportions (GIA and LES populations of western Greece form a single group). The maximum value of five clusters further partitions the Sicilian population (SIC) as a separate group. It is apparent that there is no substructure in the Central and Northern Greek population, CNG (Fig. 2).

Because of the lack of any structure among the subpopulations of Central Greece (TRI, LAR, VOL) and Kilis (KIL), we pooled the individuals from KIL into the central Greek group (CGR) to estimate the directional relative migration networks using divMigrate. The migration network (Fig. 3) reflects at some level the genetic structuring revealed by STRUCTURE analysis, as the core populations of the region (APU, CGR) cluster closely in the network space showing high gene flow and relatively low differentiation between them. The GIA and KAL populations also exhibited a relatively high gene flow with the core populations, with the former (GIA) showing connection with both APU and CGR while the latter (KAL) presents high gene flow rates only with CGR. The rest of the groups showed relatively reduced gene flow towards the core populations. It should be noted that as a general pattern, all peripheral populations seem to exchange migrants exclusively with the core populations but not between them, appearing isolated from each other (i.e. a star-shaped pattern of the migration network; Fig. 3), resembling a mainland-island metapopulation type (and not a patchy population type). However, there was no evidence of significantly asymmetric gene flow between any pair of populations.

DEMOGRAPHIC PARAMETERS

M ratios were found to be lower than the threshold value of 0.68 in all populations, which according to Garza & Williamson (2001) suggests that all populations have suffered a past

bottleneck event. The Wilcoxon signed rank test conducted in BOTTLENECK software, detected signs of a recent population bottleneck for Sicily (SIC), Limnos island (LIM) and Israel (ISR) populations (Table 3). On the other hand, inspection of plots of allele frequency classes for recent bottlenecks did not reveal any mode-shift distortion in any of the populations. However, they did show that the Croatian population (CRO) is moving towards allele fixation, since this was the only population that exhibited an allele frequency in the class 0.9-1.0 (see Supporting information, Figure S4). Most of the peripheral populations were found to have small effective population sizes (Table 3), ranging from 24.2 (LIM) to 88.4 (LES). However, in some cases (SIC, LES and ISR populations) results should be treated with caution as 95% CIs were broad.

DISCUSSION

Across all sampled populations, STRUCTURE analysis suggested the presence of two major clusters concurring with the longitudinal distribution of the species; a ‘western’ (European) and an ‘eastern’ (Asian) cluster that are both represented in the population of Israel. This result reflects the proposed phylogeographic pattern of the species; based on the mitochondrial Cyt b region, European and Asian populations were found to be divergent whereas birds from Israel seem to cluster with both of them indicating a degree of genetic mixing (Wink *et al.*, 2004). Interestingly, a comparison of plumage patterns of Lesser Kestrels unveiled substantial differences between individuals of European and Chinese descent whereas birds from Asian/Middle East populations (including Israel) exhibit an intermediate plumage pattern (Corso *et al.*, 2016). In addition, this eastern cluster is highly represented in the eastern Greek population of Limnos Island (LIM) and extends gradually up to central Greece suggesting a population consisting of birds of eastern origin possibly from populations of Asia Minor. Future inclusion of samples from Western Anatolia (i.e. Turkey) and the Middle East could shed light on such hypotheses.

Despite the high private allelic richness, suggesting that gene flow could be restricted (at some level), our analysis showed low F_{ST} values among the core populations of Europe (Spain, Apulia region in Italy and Central Greece) which is in concordance with previous larger-scale studies (Alcaide *et al.*, 2009). D_{est} on the other hand, generally showed higher values of differentiation among populations including the core ones (ES and APU). We found significant levels of differentiation among CE Mediterranean populations and a correlation of pairwise genetic and geographic distance. We should note that STRUCTURE may overestimate genetic structure in datasets characterized by such correlation (Frantz *et al.*, 2009). Correlation between genetic and geographic distance has been previously reported for the species at both local (Ortego *et al.*, 2008b) and large spatial scales (Alcaide *et al.*, 2008a; Alcaide *et al.*, 2008b) and can be driven by the distance-dependent dispersal exhibited by the species (Serrano & Tella, 2003; Serrano *et al.*, 2003; Ortego *et al.*, 2008b). Some long-distance dispersal events that could have facilitated gene flow among populations have been reported across the CE Mediterranean (Gustin, Mendi & Pedrelli, 2011; Bounas *et al.* 2016a). Nevertheless, some of these movements took place in the '50s when the species showed a wider distribution. Actually, restricted dispersal in a fragmented range, coupled with high philopatry rates, along with a relatively short generation time of the species (average lifespan of 4-6 years; Newton & Olsen, 1990; Negro, 1997), might have contributed to a more frequent individual turnover in the breeding colonies thus allowing relatively quick changes in allele frequencies that led to the patterns of genetic differentiation observed in the region (Ortego *et al.*, 2008a).

Our results for genetic bottlenecks seem to be contrasting, at first glance. M ratios suggest that the reported past population declines have indeed left genetic bottleneck signs in all populations throughout the species' range while tests for heterozygosity excess suggested bottlenecks only in three populations. Detection of a bottleneck using M ratios but not heterozygosity excess is expected when a bottleneck is older, more severe, and/or

the population has recovered (Williamson-Natesan, 2005). Thus Lesser Kestrel populations have at some point in the past undergone a severe, prolonged bottleneck, whereas in some areas (ISR, SIC, LIM) they seem to have experienced more recent, population declines. The Israeli population (ISR) has gone through a steep decline and is estimated to be less than 10% of the population prior to 1950 (Liven-Schulman *et al.*, 2004). The geographic position of Israel population could explain the increased levels of diversity despite its small N_E as a result of the genetic admixture of immigrants from other European or Asian populations in the area. Recent bottleneck signs were also detected in two of the peripheral populations in the CE Mediterranean region (SIC, LIM), that were also differentiated and showed a relatively limited gene flow with other populations (Fig. 3) and small N_E but nevertheless similar diversity patterns. Despite its small N_E , Ioannina (GIA) population exhibits high levels of diversity and gene flow with both core populations of CE Mediterranean (APU, CNG), a process that can alleviate the bottleneck effects. Since Ioannina is a known premigratory site for the species (Bounas *et al.*, 2016b), these high levels of diversity could be explained by immigrants from other populations, that is individuals that visit the site during premigration and return to breed there. Besides, it has been suggested that non-breeding distributions can shape the genetic structure of populations (Szczyś, Oswald & Arnold, 2017).

Finally, genetic drift could also play a role on the observed genetic patterns of populations in the region, since they exhibit small size and limited gene flow with other populations. This is particularly evident in the small Croatian population. This population consists of only 25 pairs and was recently discovered (Mikulic *et al.*, 2013). Therefore we are ignorant regarding its history, that is if the population was recently founded or present historically but not detected. The population did not show any signs of bottleneck but there seems to be a loss of rare alleles as well as signs of allele fixation (Table 1, Supporting information, Figure S4). Despite that, there is no observed decrease in heterozygosity, which could be consistent with a scenario of a recently founded population that experienced the

effects of genetic drift, since heterozygosity can be relatively insensitive to the loss of rare alleles due to drift (Allendorf, 1986; Allendorf, Aitken & Luikart, 2013).

CONSERVATION IMPLICATIONS

According to our results, the genetic structure of the CE Mediterranean populations of the Lesser Kestrel appears to reflect to some extent the demographic decline that led to its range fragmentation. Therefore, we suggest the management of the whole CE Mediterranean population as a single conservation unit (ESU). However, the population could be divided into four different management units (MUs) since three peripheral populations (CRO, LIM, SIC) showed limited migration rates and different allele frequencies from all the other populations. Thus a mixed strategy that aims to preserve the high diversity of the core populations of Italy and Greece, while focusing on the recovery of bottlenecked peripheral populations (SIC, LIM), should be appropriate for the conservation of the species in the region.

Translocations of individuals from the core populations to the peripheral or newly founded colonies could be used as an effective conservation action (Morandini *et al.*, 2017). Based solely on our results from the microsatellite analysis, the use of birds from the core populations seems to be an acceptable action since they were found to exhibit high diversity and low differentiation from all the others. However, the fact that our results were based solely on the analysis of selectively neutral loci prevents us from conclusions regarding the adaptive and evolutionary consequences of such action (Holderegger, Kamm & Gugerli, 2006). Thus any future translocation programmes should consider maximizing both the genetic and the adaptive similarity between populations.

ACKNOWLEDGEMENTS

We are grateful to Action for Wildlife and ANIMA rehabilitation centers, the Lesser Kestrel Recovery project team (LIFE11 NAT/BG/360), J. Hernandez-Pliego and G. Giglio for helping with sample collection. D. Vavylis, G. Vakis and E. Toli assisted in the fieldwork. LIFE for the Lesser Kestrel project team (LIFE+ 11 NAT/GR/1011), and namely K. Vlachopoulos, S. Polymeros and A. Evangelidis greatly helped with sampling in Central Greece. N. Chakarov and M. Stubbe kindly provided samples from Mongolia and we are thankful for that. V. Saravia, P. Kordopatis and S. Marin provided valuable comments on a previous draft of this manuscript. We are grateful to J.A. Allen and two anonymous reviewers for their comments that greatly improved this manuscript. This study was financially supported by a scholarship awarded to A.B. by the A.G. Leventis Foundation.

REFERENCES

- Alberto F. 2009.** MsatAllele_1. 0: an R package to visualize the binning of microsatellite alleles. *Journal of Heredity* **100**: 394-397.
- Alcaide M, Edwards SV, Negro JJ, Serrano D, Tella JL. 2008a.** Extensive polymorphism and geographical variation at a positively selected MHC class II B gene of the lesser kestrel (*Falco naumanni*). *Molecular Ecology* **17**: 2652-2665.
- Alcaide M, Serrano D, Negro JJ, Tella JL, Laaksonen T, Müller C, Gal A, Korpimäki E. 2008b.** Population fragmentation leads to isolation by distance but not genetic impoverishment in the philopatric Lesser Kestrel: a comparison with the widespread and sympatric Eurasian Kestrel. *Heredity* **102**: 190-198.
- Alcaide M, Serrano D, Tella JL, Negro JJ. 2009.** Strong philopatry derived from capture–recapture records does not lead to fine-scale genetic differentiation in Lesser Kestrels. *Journal of Animal Ecology* **78**: 468-475.
- Alcala N, Goudet J, Vuilleumier S. 2014.** On the transition of genetic differentiation from isolation to panmixia: what we can learn from GST and D. *Theoretical population biology* **93**: 75-84.
- Allendorf FW. 1986.** Genetic drift and the loss of alleles versus heterozygosity. *Zoo Biology* **5**: 181-190.
- Allendorf FW, Aitken SN, Luikart G. 2013.** *Conservation and the genetics of populations*. 2nd edn. Wiley-Blackwell Publishing, Oxford.
- Amos W, Balmford A. 2001.** When does conservation genetics matter? *Heredity* **87**: 257-265.
- Benjamini Y, Hochberg Y. 1995.** Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the royal statistical society* **57**: 289-300.
- BirdLife International. 2017a.** European birds of conservation concern: populations, trends and national responsibilities. BirdLife International, Cambridge.

- BirdLife International. 2017b.** Species factsheet: *Falco naumanni*. Downloaded from <http://www.birdlife.org> on 21/03/2017.
- Bounas A, Panuccio M, Evangelidis A, Sotiropoulos K, Barboutis C. 2016a.** The Migration of the Lesser Kestrel *Falco naumanni* in Eastern Europe-A Ringing Recovery and Direct Observation Approach. *Acrocephalus* **37**: 49-56.
- Bounas A, Tsiakiris R, Vlachopoulos K, Bukas N, Stara K, Sotiropoulos K. 2016b.** Large Premigratory Roost of Lesser Kestrels (*Falco naumanni*) in Ioannina City, Greece: Trends, Roost Characteristics, and Implications for Conservation. *Journal of Raptor Research* **50**: 416-421.
- Cornuet JM, Luikart G. 1996.** Description and power analysis of two tests for detecting recent population bottlenecks from allele frequency data. *Genetics* **144**: 2001-2014.
- Corso A, Vigano M, Jansen JJ, Starnini L. 2016.** Geographical plumage variation in Lesser Kestrel. *Dutch Birding* **38**: 271-292.
- Cramp S, Simmons K. 1980.** Birds of the Western Palearctic, Vol. 2: Oxford University Press, Oxford.
- Cristescu R, Sherwin WB, Handasyde K, Cahill V, Cooper DW. 2010.** Detecting bottlenecks using BOTTLENECK 1.2. 02 in wild populations: the importance of the microsatellite structure. *Conservation Genetics* **11**: 1043-1049.
- Di Maggio R, Mengoni C, Mucci N, Campobello D, Randi E, Sarà M. 2014.** Do not disturb the family: roles of colony size and human disturbance in the genetic structure of lesser kestrel. *Journal of Zoology* **295**: 108-115.
- Do C, Waples RS, Peel D, Macbeth G, Tillett BJ, Ovenden JR. 2014.** NeEstimator v2: re-implementation of software for the estimation of contemporary effective population size (N_e) from genetic data. *Molecular Ecology Resources* **14**: 209-214.
- Dyer R. 2014.** GSTUDIO: Analyses and functions related to the spatial analysis of genetic marker data. *R package version 3*.
- Earl DA, vonHoldt BM. 2012.** STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genetics Resources* **4**: 359-361.
- Evanno G, Regnaut S, Goudet J. 2005.** Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology* **14**: 2611-2620.
- Excoffier L, Lischer HE. 2010.** Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources* **10**: 564-567.
- Falush D, Stephens M, Pritchard JK. 2003.** Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics* **164**: 1567-1587.
- Frantz A, Cellina S, Krier A, Schley L, Burke T. 2009.** Using spatial Bayesian methods to determine the genetic structure of a continuously distributed population: clusters or isolation by distance? *Journal of Applied Ecology* **46**: 493-505.
- Garcia JT, Alda F, Terraube J, Mougeot F, Sternalski A, Bretagnolle V, Arroyo B. 2011.** Demographic history, genetic structure and gene flow in a steppe-associated raptor species. *BMC Evolutionary Biology* **11**: 333.
- Gerlach G, Jueterbock A, Kraemer P, Deppermann J, Harmand P. 2010.** Calculations of population differentiation based on GST and D: forget GST but not all of statistics! *Molecular Ecology* **19**: 3845-3852.
- Goudet J. 2002.** FSTAT version 2.9. 3.2, a program to estimate and test gene diversities and fixation indices. *Institute of Ecology, Lausanne, Switzerland. Website <http://www2.unil.ch/popgen/softwares/fstat.htm> [accessed 20 August 2012]*.
- Gustin M, Mendi M, Pedrelli M. 2011.** Grillaio nato a Montescaglioso (MT) si riproduce in Emilia-Romagna. *Picus* **37**: 83 (in Italian).

487 **Hanski I, Gaggiotti OE. 2004.** *Ecology, genetics, and evolution of metapopulations*. Academic
488 Press.

489 **Hill WG. 1981.** Estimation of effective population size from data on linkage disequilibrium.
490 *Genetical research* **38**: 209-216.

491 **Holderegger R, Kamm U, Gugerli F. 2006.** Adaptive vs. neutral genetic diversity: implications
492 for landscape genetics. *Landscape Ecology* **21**: 797-807.

493 **Hubisz MJ, Falush D, Stephens M, Pritchard JK. 2009.** Inferring weak population structure
494 with the assistance of sample group information. *Molecular Ecology Resources* **9**:
495 1322-1332.

496 **Iñigo A, Barov B. 2010.** Action Plan for the Lesser Kestrel (*Falco naumanni*) in the European
497 Union. SEO Birdlife & Birdlife International for the European Commission, Madrid.

498 **Jensen JL, Bohonak AJ, Kelley ST. 2005.** Isolation by distance, web service. *BMC genetics* **6**:
499 13.

500 **Kalinowski ST. 2005.** HP-Rare: a computer program for performing rarefaction on measures
501 of allelic diversity. *Molecular Ecology Notes* **5**: 187-189.

502 **Keenan K, McGinnity P, Cross TF, Crozier WW, Prodöhl PA. 2013.** diveRsity: an R package
503 for the estimation and exploration of population genetics parameters and their
504 associated errors. *Methods in Ecology and Evolution* **4**: 782-788.

505 **Kmetova E, Zhelev P, Mechev A, Gradev G, Ivanov I. 2012.** Natural Colonies of Lesser
506 Kestrel (*Falco naumanni*) in European Turkey and Discussion on the Chances of
507 Natural Re-colonization of the Species in Bulgaria. *Acta Zoologica Bulgarica* **4**: 47-54.

508 **Lagnel J. 2015.** ParaStructure. GitHub repository [https://github.com/jacqueslagnel/](https://github.com/jacqueslagnel/parastructure)
509 [parastructure](https://github.com/jacqueslagnel/parastructure).

510 **Liven-Schulman I, Leshem Y, Alon D, Yom-Tov Y. 2004.** Causes of population declines of the
511 Lesser Kestrel *Falco naumanni* in Israel. *Ibis* **146**: 145-152.

512 **Luikart G, Allendorf F, Cornuet J, Sherwin W. 1998.** Distortion of allele frequency
513 distributions provides a test for recent population bottlenecks. *Journal of Heredity*
514 **89**: 238-247.

515 **Meirmans PG. 2015.** Seven common mistakes in population genetics and how to avoid
516 them. *Molecular Ecology* **24**: 3223-3231.

517 **Mikulic K, Ivan Budinski, Antica Culina, Luka Jurinovic, Lucic V. 2013.** The return of the
518 Lesser Kestrel (*Falco naumanni*) as a breeding bird to Croatia. *Acrocephalus* **34**: 71-
519 74.

520 **Miller MP, Mullins TD, Parrish Jr JW, Walters JR, Haig SM. 2012.** Variation in migratory
521 behavior influences regional genetic diversity and structure among American Kestrel
522 populations (*Falco sparverius*) in North America. *Journal of Heredity* **103**: 503-514.

523 **Morandini V, Benito E, Newton I, Ferrer M. 2017.** Natural expansion versus translocation in
524 a previously human-persecuted bird of prey. *Ecology and Evolution* **7**: 3682-3688.

525 **Moritz C. 1994.** Defining 'evolutionarily significant units' for conservation. *Trends in Ecology*
526 *& Evolution* **9**: 373-375.

527 **Negro JJ. 1997.** Lesser Kestrel *Falco naumanni*. Birds of the Western Palearctic. Update 1,
528 49-56..

529 **Negro JJ, Hiraldo F, Donazar JA. 1997.** Causes of natal dispersal in the lesser kestrel:
530 inbreeding avoidance or resource competition? *Journal of Animal Ecology* **66**: 640-
531 648.

532 **Nesje M, Røed KH, Lifjeld JT, Lindberg P, Steen OF. 2000.** Genetic relationships in the
533 peregrine falcon (*Falco peregrinus*) analysed by microsatellite DNA markers.
534 *Molecular Ecology* **9**: 53-60.

535 **Newton I, Olsen P. 1990.** *Birds of prey*. Merehurst Press, London.

Ortego J, Aparicio JM, Cordero PJ, Calabuig G. 2008a. Individual genetic diversity correlates with the size and spatial isolation of natal colonies in a bird metapopulation. *Proceedings of the Royal Society of London B: Biological Sciences* **275**: 2039-2047.

Ortego J, Calabuig G, Aparicio JM, Cordero PJ. 2008b. Genetic consequences of natal dispersal in the colonial lesser kestrel. *Molecular Ecology* **17**: 2051-2059.

Ortego J, González EG, Sánchez-Barbudo I, Aparicio JM, Cordero PJ. 2007. Novel highly polymorphic loci and cross-amplified microsatellites for the lesser kestrel *Falco naumanni*. *Ardeola* **54**: 101-108.

Padilla JA, Parejo JC, Salazar J, Martínez-Trancón M, Rabasco A, Sansinforiano E, Quesada A. 2009. Isolation and characterization of polymorphic microsatellite markers in lesser kestrel (*Falco naumanni*) and cross-amplification in common kestrel (*Falco tinnunculus*). *Conservation Genetics* **10**: 1357-1360.

Palsbøll PJ, Berube M, Allendorf FW. 2007. Identification of management units using population genetic data. *Trends in Ecology & Evolution* **22**: 11-16.

Peakall R, Smouse PE. 2012. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics* **28**: 2537-2539.

Piry S, Luikart G, Cornuet JM. 1999. BOTTLENECK: a computer program for detecting recent reductions in the effective size using allele frequency data. *Journal of Heredity* **90**: 502-503.

Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure using multilocus genotype data. *Genetics* **155**: 945-959.

R Core Team. 2015. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at: <https://www.R-project.org/>.

Rodriguez A, Negro JJ, Bustamante J, Fox JW, Afanasyev V. 2009. Geolocators map the wintering grounds of threatened Lesser Kestrels in Africa. *Diversity and Distributions* **15**: 1010-1016.

Rosenberg NA. 2004. DISTRUCT: a program for the graphical display of population structure. *Molecular Ecology Notes* **4**: 137-138.

Sarà M. 2010. Climate and land-use changes as determinants of lesser kestrel *Falco naumanni* abundance in Mediterranean cereal steppes (Sicily). *Ardeola* **57**: 3-22.

Serrano D, Tella JL. 2003. Dispersal within a spatially structured population of lesser kestrels: the role of spatial isolation and conspecific attraction. *Journal of Animal Ecology* **72**: 400-410.

Serrano D, Tella JL, Donazar JA, Pomarol M. 2003. Social and individual features affecting natal dispersal in the colonial lesser kestrel. *Ecology* **84**: 3044-3054.

Sundqvist L, Keenan K, Zackrisson M, Prodöhl P, Kleinhans D. 2016. Directional genetic differentiation and relative migration. *Ecology and Evolution* **6**: 3461-3475.

Szczys P, Oswald SA, Arnold JM. 2017. Conservation implications of long-distance migration routes: Regional metapopulation structure, asymmetrical dispersal, and population declines. *Biological Conservation* **209**: 263-272.

Taylor B, Dizon A. 1999. First policy then science: why a management unit based solely on genetic criteria cannot work. *Molecular Ecology* **8**.

Toonen RJ, Hughes S. 2001. Increased throughput for fragment analysis on an ABI Prism® 377 automated sequencer using a membrane comb and STRand software. *Biotechniques* **31**: 1320-1325.

Uzunova D, Lisichanets E. 2016. Lesser Kestrel's population and conservation status in Macedonia. In: *Abstracts of the International Lesser Kestrel Expert Workshop*, Plovdiv, Bulgaria.

Van Oosterhout C, Hutchinson WF, Wills DP, Shipley P. 2004. MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**: 535-538.

- Vucetich JA, Waite TA. 2003.** Spatial patterns of demography and genetic processes across the species' range: null hypotheses for landscape conservation genetics. *Conservation Genetics* **4**: 639-645.
- Waples RS. 2006.** A bias correction for estimates of effective population size based on linkage disequilibrium at unlinked gene loci. *Conservation Genetics* **7**: 167.
- Waples RS, Do C. 2010.** Linkage disequilibrium estimates of contemporary Ne using highly variable genetic markers: a largely untapped resource for applied conservation and evolution. *Evolutionary Applications* **3**: 244-262.
- Williamson-Natesan EG. 2005.** Comparison of methods for detecting bottlenecks from microsatellite loci. *Conservation Genetics* **6**: 551-562.
- Willoughby JR, Sundaram M, Wijayawardena BK, Lamb MC, Kimble SJ, Ji Y, Fernandez NB, Antonides JD, Marra NJ, Dewoody JA. 2017.** Biome and migratory behaviour significantly influence vertebrate genetic diversity. *Biological Journal of the Linnean Society* **121**: 446-457.
- Wink M, Sauer-Gurth H, Pepler D. 2004.** Phylogeographic relationships of the Lesser Kestrel (*Falco naumanni*) in breeding and wintering quarters inferred from nucleotide sequences of the mitochondrial cytochrome b gene. *Raptors worldwide*, 505-510.

617

618 **Figure 1.** Map of the Lesser Kestrel populations used in the present study. SES: Andalucia,
 619 ES/BG: Extremadura, APU: Apulia, SIC: Sicily, CRO: Croatia, GIA: Ioannina, LES: Agrinio, TRI:
 620 Trikala, LAR: Larisa, VOL: Volos, KIL: Kilis, KAL: Komotini, LIM: Limnos, ISR: Israel, MON:
 621 Mongolia. Shaded areas represent the breeding distribution of the Lesser Kestrel (modified
 622 from BirdLife International). Sampling sites pooled for the analyses are circled (ES: Spain;
 623 CNG: Central-North Greece)

624 **Figure 2.** Admixture proportions (proportions of membership to each of K inferred clusters) of
 625 individual Lesser Kestrels. Upper plots correspond to the number of clusters when the full
 626 dataset was modeled (K = 2, K = 3), whereas the lower 3 plots show admixture proportions of
 627 the CE Mediterranean populations only, for K=2, K=4 and K=5 inferred clusters respectively

628 **Figure 3.** Directional relative migration networks of grouped Lesser Kestrel populations from
 629 CE Mediterranean. Networks were visualized with divMigrate using Alcalá's Nm. (a) all relative
 630 migration rate values (b) only values above 0.2 threshold (c) only gene flow values estimated
 631 >0.4. Line shading and thickness increases with the relative strength of gene flow.

632

633

634

635

Table 1 Measures of genetic variation of all sampled Lesser Kestrel populations. Number of genotyped individuals (N), number of alleles per locus (A), allelic richness (A_R), observed (H_O) and expected (H_E) heterozygosity, private allelic richness (π) and inbreeding coefficient (F_{IS}). Values are presented as means $\pm$ standard error (in brackets)

Code	Location	N	A	A_R	H_O	H_E	π	F_{IS}
SES	Andalucia	19	7.1 (0.9)	5.4 (0.6)	0.635 (0.037)	0.696 (0.037)	0.17	0.082 (0.041)
ES/BG	Extremadura	25	8.3 (1.3)	5.6 (0.7)	0.674 (0.040)	0.699 (0.037)	0.26	0.029 (0.037)
APU	Apulia	44	9.4 (1.5)	5.5 (0.6)	0.669 (0.038)	0.707 (0.036)	0.13	0.053 (0.025)
SIC	Sicily	12	5.9 (0.8)	5.1 (0.6)	0.682 (0.038)	0.667 (0.041)	0.18	-0.045 (0.045)
CRO	Croatia	14	5.6 (0.6)	4.7 (0.5)	0.654 (0.048)	0.650 (0.048)	0.03	-0.023 (0.043)
GIA	Ioannina	24	7.6 (1.1)	5.2 (0.6)	0.634 (0.051)	0.654 (0.048)	0.16	0.036 (0.031)
LES	Agrinio	16	6.7 (0.8)	5.4 (0.6)	0.603 (0.058)	0.684 (0.042)	0.18	0.136 (0.047)
TRI	Trikala	20	7.6 (1.3)	5.5 (0.7)	0.626 (0.056)	0.680 (0.046)	0.08	0.087 (0.046)
LAR	Larisa	20	7.4 (1.1)	5.4 (0.6)	0.669 (0.053)	0.680 (0.044)	0.09	0.024 (0.045)
VOL	Volos	20	7.6 (1.2)	5.5 (0.7)	0.678 (0.051)	0.662 (0.048)	0.13	-0.027 (0.035)
KIL	Kilkis	13	6.4 (0.9)	5.3 (0.7)	0.621 (0.062)	0.661 (0.048)	0.09	0.067 (0.064)
KAL	Komotini	20	7.1 (1.1)	5.3 (0.6)	0.689 (0.053)	0.677 (0.046)	0.04	-0.019 (0.040)
LIM	Limnos	11	5.1 (0.6)	4.8 (0.5)	0.661 (0.060)	0.667 (0.041)	0.09	0.024 (0.060)
ISR	Israel	20	7.3 (1.1)	5.5 (0.7)	0.619 (0.062)	0.682 (0.042)	0.28	0.124 (0.055)
MON	Mongolia	17	7.2 (1.0)	5.4 (0.6)	0.585 (0.057)	0.655 (0.049)	0.52	0.155 (0.046)

643 **Table 2** Pairwise F_{st} -values (below diagonal) and D_{est} -values (above diagonal) among Lesser Kestrel populations. Statistically significant values after B-H
644 correction for multiple comparisons are given in bold ($p < 0.039$)

645		SES	ES/BG	APU	SIC	CRO	GIA	LES	TRI	LAR	VOL	KIL	KAL	LIM	ISR	MON
646	SES	-	0.018	0.038	0.099	0.132	0.048	0.053	0.014	0.030	0.053	0.016	0.027	0.147	0.059	0.093
	ES/BG	0.000	-	0.035	0.096	0.138	0.056	0.049	0.015	0.046	0.003	0.029	0.041	0.160	0.069	0.126
647	APU	0.001	0.004	-	0.100	0.103	0.076	0.072	0.014	0.021	0.041	0.030	0.052	0.143	0.054	0.121
	SIC	0.026	0.035	0.029	-	0.167	0.105	0.128	0.111	0.099	0.108	0.069	0.096	0.175	0.115	0.183
648	CRO	0.033	0.032	0.026	0.048	-	0.152	0.117	0.101	0.112	0.146	0.117	0.130	0.191	0.117	0.168
	GIA	0.012	0.018	0.019	0.035	0.044	-	0.077	0.027	0.078	0.065	0.026	0.054	0.143	0.088	0.151
649	LES	0.013	0.014	0.019	0.038	0.033	0.029	-	0.065	0.084	0.072	0.050	0.067	0.183	0.074	0.158
650	TRI	0.003	0.005	0.004	0.033	0.027	0.009	0.023	-	0.000	0.035	0.000	0.000	0.100	0.030	0.110
	LAR	0.005	0.010	0.006	0.032	0.036	0.019	0.030	0.000	-	0.027	0.005	0.028	0.153	0.064	0.112
651	VOL	0.010	0.004	0.013	0.030	0.037	0.018	0.018	0.012	0.012	-	0.015	0.047	0.132	0.080	0.121
	KIL	0.001	0.009	0.008	0.016	0.034	0.004	0.018	0.000	0.000	0.004	-	0.037	0.136	0.020	0.095
652	KAL	0.011	0.013	0.021	0.038	0.044	0.025	0.024	0.005	0.010	0.021	0.019	-	0.092	0.088	0.085
653	LIM	0.035	0.035	0.036	0.051	0.064	0.041	0.046	0.020	0.034	0.040	0.039	0.019	-	0.152	0.143
	ISR	0.017	0.019	0.015	0.029	0.035	0.029	0.022	0.015	0.026	0.024	0.012	0.035	0.045	-	0.094
654	MON	0.032	0.035	0.042	0.055	0.060	0.052	0.053	0.038	0.039	0.043	0.039	0.024	0.042	0.033	-

Table 3. Contemporary effective population sizes (N_E) and 95% CI, results of tests for genetic bottlenecks: Garza-Williamson M values, Wilcoxon sign-rank tests for heterozygosity excess and mode-shift tests for all Lesser Kestrel populations. The Wilcoxon tests were carried using the TPM model (5% SMM, 95% IAM). Values in bold are indicative of a bottleneck ($M < 0.68$ for the Garza-Williamson ratio and $P < 0.05$ for the Wilcoxon test)

Code	Location	N_E	M	P (TPM)	Allele frequency distribution
ES	Spain	Not estimated	0.43	0.083	L-shaped
APU	Apulia	Not estimated	0.45	0.148	L-shaped
SIC	Sicily	57.3 (29.8-327.7)	0.39	0.007	L-shaped
CRO	Croatia	47.7 (21.6-82.8)	0.41	0.216	L-shaped
GIA	Ioannina	26.6 (19.5-39.1)	0.45	0.390	L-shaped
LES	Agrinio	88.4 (40.5-167.8)	0.45	0.056	L-shaped
CNG	Central-North Greece	Not estimated	0.45	0.078	L-shaped
LIM	Limnos	24.2 (14.5-55.8)	0.42	0.004	L-shaped
ISR	Israel	54.9 (35.8-106.1)	0.43	0.009	L-shaped
MON	Mongolia	Not estimated	0.42	0.201	L-shaped

672



673

674 **Figure 1**




Figure 2




Figure 3